Diabetes & the Medicare Population: Idaho

Introduction to the Report

Diabetes has a considerable impact on the nation's healthcare system and on individuals living with the condition. This report is intended to support efforts within Washington's communities to monitor and improve diabetes-related quality measures. Data represents the year of July 2016 – June 2017. This report's population also includes individuals under age 65 who qualify for Medicare due to chronic disability (approximately 20% of the Medicare population). It does not include data on individuals living outside the state who may seek care in the state.

Through our work as Medicare's Quality Improvement Network – Quality Improvement Organization (QIN-QIO) in Idaho and Washington, Qualis Health is assisting providers and patients to develop systems to better manage and monitor this disease. We are also working with communities throughout both states to improve care coordination, prevent adverse drug events, and reduce unnecessary rehospitalizations. To learn more about our QIN-QIO activities, please see:

- Diabetes-related consulting for primary care physicians www.Medicare.QualisHealth.org/DiabetesProvider
- Diabetes self-management education for Medicare beneficiaries and their family members and caregivers www.Medicare.QualisHealth.org/EDC
- Consulting related to care coordination and reducing unnecessary rehospitalizations on a community level
 www.Medicare.QualisHealth.org/Transitions





This material was prepared by Qualis Health, the Medicare Quality Innovation Network - Quality Improvement Organization (QIN-QIO) for Idaho and Washington, under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. The contents presented do not necessarily reflect CMS policy. ID-C3-QH-3249-11-17

Disease Prevalence

Valley

Over 40,000 of Idaho's Medicare beneficiaries have been diagnosed with diabetes. While diabetes is prevalent throughout the state, the difference between the communities with the lowest and highest rates is stark: the rate of diabetes among Medicare beneficiaries residing in Eastern Idaho is more than 40% higher than that of Treasure Valley.

30% 26.7% 25.3% 23.4% 25% 22.3% 21.9% 19.4% 20% 18.3% 15% 10% 5% 0% Treasure Northern Idaho North Central Magic Valley Greater Southeast Eastern Idaho

Figure 1: Prevalence of Diabetes among Medicare Beneficiaries by Community, July 2016 – June 2017


Medicare beneficiaries with diabetes also face increased morbidity from other chronic conditions. Figure 2 shows the prevalence for select chronic conditions for beneficiaries with and without diabetes.


State Rate (21.4%)

Canyon

County


Idaho





There are also racial disparities in the prevalence of diabetes, as shown in Figure 3 below.

Figure 3: Prevalence of Diabetes among Medicare Beneficiaries by Race and Dual-Eligibility with Medicaid, July 2016 – June 2017



Appropriateness of Care

People with diabetes have greatly increased risk for cardiovascular disease and other comorbidities, so screening for glycemic and lipid control and neuropathic complications is part of ongoing care management. The charts below show variation by community for recommended care measures.

Figure 4: Percent of Medicare Beneficiaries with Diabetes Receiving Annual HbA1c Test by Community, July 2016 – June 2017




Figure 5: Percent of Medicare Beneficiaries with Diabetes Receiving Annual Lipid Test by Community, July 2016 – June 2017




Figure 6: Percent of Medicare Beneficiaries with Diabetes Receiving Annual Eye Exam by Community, July 2016 – June 2017




Figure 7: Percent of Medicare Beneficiaries with Diabetes Ever Receiving Pneumococcal Vaccine by Community, July 2016 – June 2017




Figure 8: Percent of Medicare Beneficiaries with Diabetes Receiving Annual Influenza Vaccination by Community, July 2016 – June 2017




Figure 9: Percent of Medicare Beneficiaries with Diabetes Ever Receiving Pneumococcal Vaccine by Rurality, July 2016 – June 2017





Figure 10: Percent of Medicare Beneficiaries with Diabetes Receiving Annual Influenza Vaccination by Rurality, July 2016 – June 2017



Hospital Utilization and Medicare Spending

Medicare beneficiaries with diabetes have increased admission, 30-day rehospitalization, and Emergency Department (ED) rates compared to beneficiaries without diabetes. However, that utilization is largely driven by individuals who have diabetes that has progressed to Chronic Kidney Disease (CKD) and End Stage Renal Disease (ESRD). The figures below show utilization, mortality, and per capita costs for individuals by diabetes status.

Figure 11: ED Visits, Hospital Admissions, and 30-Day Re-Hospitalizations by Diabetes Status for Medicare Beneficiaries, July 2016 – June 2017






Figure 12: Mortality by Diabetes Status for Medicare Beneficiaries, July 2016 – June 2017

Figure 13: Per Capita Costs by Diabetes Status for Medicare Beneficiaries, July 2016 – June 2017





In part as a result of racial disparities in diabetes progression, there are racial disparities in utilization among beneficiaries with diabetes.

Figure 14: Admits per 1,000 Beneficiaries with Diabetes by Race, July 2016 – June 2017




Figure 15: ED Visits per 1,000 Beneficiaries with Diabetes by Race, July 2016 – June 2017






Figure 16: 30-Day Rehospitalizations per 1,000 Beneficiaries with Diabetes by Race, July 2016 – June 2017


Figure 17: LEAs per 100,000 Beneficiaries by Race, July 2016 – June 2017





Diabetes Self-Management Education Outcomes

Qualis Health is currently partnering with community organizations to support Diabetes Self-Management Education by offering the evidence-based Diabetes Self-Management Workshops. This gold-standard strategy, originally developed at Stanford University, empowers people with diabetes to manage their symptoms, address day-to-day problems and improve the quality of their lives. Participants learn, practice and master skills such as goal-setting and action -planning, blood sugar monitoring, medication management, meal planning, safe physical activity, stress and depression management, problem-solving and decision-making.



Pre- and post-class patient activation surveys indicate that participants¹

- Participants were more likely to feel able to ask their doctor questions about their treatment plans after participation (68% before vs. 94% after).
- Participants were more likely to feel able to make a plan with goals that will help control their diabetes after participation (55% before versus 91% after).

¹ Complete national and state results of pre-class and post-class Patient Activation Surveys are available upon request. Organizations partnering with Qualis Health can receive this data for participants of their specific classes in aggregate form after they have had at least 20 individuals complete pre- and post-class surveys.

Community Data Tables

Table 1: Utilization and Per Capita Costs for Medicare Beneficiaries with Diabetes by Community, July 2016 – June 2017

	Percent with CKD	ED Visits per 1,000	Admissions per 1,000	30-Day Rehospitalizations per 1,000	Lower Extremity Amputations per 100,000	Per Capita Costs
Eastern Idaho	44.9%	543.08	318.92	40.68	26.81	\$12,065
Greater Canyon County	52.6%	831.93	418.66	65.19	84.06	\$12,105
Magic Valley	49.8%	562.33	443.82	72.12	80.21	\$12,399
North Central	38.8%	859.27	378.51	54.41	52.28	\$11,945
Northern Idaho	43.9%	623.09	346.66	47.90	35.37	\$11,500
Southeast Idaho	43.7%	772.84	372.99	59.64	37.48	\$12,696
Treasure Valley	48.8%	712.81	366.80	54.46	61.09	\$11,316
Statewide	45.8%	690.14	367.08	54.23	50.23	\$11,890